Biokemistri
Vol. 32, No. 1, March 31, 2020, pages 1 - 10
Printed in Nigeria

0795-8080/2020 $10.00 + 0.00

© 2020 Nigerian Society for Experimental Biology
http://www.niseb.org.ng/journals
Also available online at http://www.bioline.org.br/bk

BKR 2020003/32101

Heavy metals correlate with cellular adenosine triphosphate and fructose levels in municipal dumpsite exposure

Oluwakemi T. Oyelowo[1]*, Oju E. Omamogho[1], Ikechukwu F. Ezenwajiaku[2], Olajumoke A. Morenikeji [2], Adekunle Mofolorunso[1]

[1]Department of Physiology, University of Lagos, Lagos, Nigeria
[2]Department of Zoology, University of Ibadan, Ibadan, Nigeria

*Author for Correspondence: Oluwakemi T. Oyelowo (ooyelowo@unilag.edu.ng) Tel: +23480367333891

(Received January 16, 2020; Accepted March 9, 2020)

ABSTRACT: This study assessed the influence of direct exposure of rodents to Awotan Solid Waste Dumpsite (ASWD) in Ibadan, Nigeria. Reproductive energy content was evaluated by assessing testicular ATP activity, testicular polychlorinated biphenyls (PCBs), testicular heavy metal and fructose levels in seminal vesicle and coagulating glands. The exposure of the male rats was from postnatal day 22 and for a period of 10 weeks. The direct municipal dumpsite exposure resulted in testicular heavy metal accumulation viz: mercury > iron > zinc > nickel > cadmium > arsenic. There was a decrease in serum lactate dehydrogenase (LDH) concentration, an increase in testicular LDH concentration (markers of cellular ATP) and a decrease in the fructose level in the seminal vesicle of the dumpsite exposed (DSE) animals compared with the control. The PCBs were not detected in the testis of DSE animals but there was significant correlation between testicular LDH and some metals. It is concluded that some possible mechanisms by which direct exposure to a dumpsite elicits energy depletion in the male rat testes could be through the inhibition of fructose level and LDH activity.

Keywords: Heavy metal, ATP, Fructose, Dumpsite, Polychlorinated biphenyls

Introduction

Reproductive disorders, high infertility rate, increased birth defects, repressed immunological function, and increased frequency of cancers have been linked to exposure to landfill sites (1,2). Landfill which refers to the deposition of waste in an area specially designed for such is a major method in waste management hierarchy in developing countries (1). Landfills may contain household or municipal wastes, industrial, biomedical or infectious wastes (3). The landfills thus can contain toxic substances such as heavy metals, dissolved organic matter, polychlorinated biphenyls and inorganic macro compounds. Some of these xenobiotics have been classified by The United States Environmental Protection Agency to be of concern to the male reproductive health (4).

Heavy metal environmental pollution even at low levels and their subsequent long-term collective health effects are among the leading causes of health concerns the world over (5). There have been studies suggesting that heavy metals are a major source of oxidative stress in the cell and that they play an important role in the aetiology of diverse human pathologies like carcinogenesis (6). Research has also revealed that heavy metals directly modify and /or damage the DNA structure (7). There have also



Fig. 2 Levels of Cd, As, Hg, and Pb in testes of rats after 10 weeks of dumpsite exposure (DSE). Each bar represents mean ± SEM of 12 rats. *Values differ significantly from the control ($p<0.05$).



Fig. 3 Fructose level in seminal vesicle and coagulating glands of rats after 10 weeks of dumpsite exposure (DSE). Each bar represents mean ± SEM of 12 rats. *Values differ significantly from the control ($p<0.05$).



Fig. 4 Serum and testis LDH levels of rats after 10 weeks of dumpsite exposure (DSE). Each bar represents mean ± SEM of 12 rats. *Values differ significantly from the control ($p<0.05$).

Seminal vesicle and coagulating gland Fructose level

There was a significant decrease in the level of fructose found in the seminal vesicle and coagulating glands compared with the control (Fig 3).

Activity of Serum and Testicular Lactate Dehydrogenase (LDH) (Functional marker of ATP).

The activity of the serum LDH was significantly inhibited in the rats from the dumpsite compared with the control rats however, the dumpsite exposure was associated with a significant increase in the testicular activities of LDH compared with the control (Fig 4). Testicular LDH significantly correlated with mercury and copper (Table 2).

Polychlorinated biphenyls (PCBs) concentration

Polychlorinated biphenyls (PCBs) were absent in the testis of the dumpsite-exposed animals.

Table 2: Correlation matrix between Heavy metals and Testicular LDH, Serum LDH, Fructose in control animals

		Cu	Ni	As	Zn	Hg	Fe	Cd	Pb
	Testicular LDH	0.66*	0.30	0.47	-0.36	0.70	0.67	0.72	0.72
Control	Serum LDH	-0.1	0.10	0.01	-0.31	0.18	0.47	-0.36	-0.36
	Fructose	-0.43	0.06	0.02	-0.58	-0.24	-0.49	-0.95	-0.95**

*Correlation is significant at the 0.05 level. **Correlation is significant at the 0.01 level.

Table 3: Correlation matrix between Heavy metals and Testicular LDH, Serum LDH, Fructose in dumpsite exposed (DSE) animals.

		Cu	Ni	As	Zn	Hg	Fe	Cd	Pb
	Testicular LDH	0.77*	-0.19	0.04	-0.60	0.83*	0.12	-0.23	0.54
DSE	Serum LDH	-0.10	-0.08	0.38	-0.22	-0.14	0.27	0.27	0.41
	Fructose	-0.55	0.01	-0.04	0.30	-0.62	-0.62	-0.62	-0.03

*Correlation is significant at the 0.05 level. **Correlation is significant at the 0.01 level.

Discussion

There was an accumulation of mercury, arsenic, cadmium, iron, zinc and nickel in the testes of the dumpsite exposed (DSE) animals. A previous study from our laboratory (28) reported that heavy metal contents found in the soil of Awotan Solid Waste Dumpsite (ASWD) which is the study area for this research were above the USEPA permissible limits which would have resulted in the accumulation of these toxic heavy metals signifying their roles in testicular toxicity observed in the DSE animals. Although the levels of lead and copper were above the USEPA permissible limits in the soil of ASWD, they were not accumulated in the testis of the animals exposed to the dumpsite. A probable reason is that metals interact additively, synergistically as well as antagonistically and affect each other's absorption, distribution and excretion. This interference with metabolism can reduce the concentration in the organism or decrease the bioavailability. The competition between lead and/or cadmium and zinc for the same binding sites in enzymes, proteins, and transporters can affect structure and/or function of cell membranes, change enzyme activity, induce oxidative stress and apoptosis which all have grave consequences on cell growth, development, and differentiation. Thus, these interactions contribute to interindividual differences in susceptibility to adverse effects of metals in men (29).

According to a study by Sharpe *et al.* (30) sertoli cells proliferate during the foetal, neonatal, and pre-pubertal periods and each of these periods is particularly sensitive to the adverse effects of metals. Testicular toxicity in rats has also been shown as a good predictor in human subjects (31). Arsenic and cadmium which are classified as the first and seventh most hazardous substances (32) were significantly increased in the DSE group. Arsenic and its compounds have been reported to disrupt ATP production when exposed to animals by way of water drinking (33). The observation in this study corroborates other studies that reported that cells exposed to arsenic showed a considerable depletion in ATP and glycogen levels in the liver and other tissues. Cadmium has been reported to cause severe damage to spermatogenic epithelium in an animal model (34). Other reports have shown that at concentrations greater than (0.003 mg/L), cadmium had direct injury on the testes by damaging germ cells and Sertoli cells and subsequently reducing sperm quality (34). The increase in cadmium level in the dumpsite-exposed animals also confirms the fact that cadmium accumulates in the male reproductive organs, especially the testis of both humans and animals (35). Cadmium has been shown to negatively affect accessory sex organ functions (10).

The increase in testicular and epididymal weights (36) has been reported in cadmium exposure which corroborates the findings of this study. The increase in the organ weights generally indicates an excessive accumulation of interstitial fluid and maybe a sensitive indicator of decreased sperm production (36). Some studies have earlier reported that cadmium mediated histological changes in the testes, epididymis and accessory sex organs (37). Moreover, exposure to cadmium, lead and arsenic may contribute to prostate cancer development and some reports have associated their exposure with increased serum prostate specific antigen (PSA) (10). Metals can affect the male reproductive system directly when they target specific reproductive organs. When there is an accumulation of metals in the testis, epididymis, prostate, seminal vesicle, they impair progressive sperm motility (38). Metals affect the secretory function of the seminal vesicles leading to the loss of fertility, libido or impotence (39).

The seminal plasma is a mixture of secretions derived principally from the major accessory sex glands such as the seminal vesicle, prostate, Cowper's glands with minor contributions from the epididymis, ampulla gland etc. Since neither the testis nor epididymal sperm contains fructose, the immotile spermatozoa encounter fructose only after it has been intermixed with seminal plasma during ejaculation. This occurs at a crucial moment when the sperm requires a high degree of motility requiring a source of quick available energy. This energy is provided by the metabolic process of fructolysis during which lactic acid is formed. Since growth and secretory function of the accessory sex glands are under the control of the testis, the level of fructose in seminal plasma appears to be a precise indicator of the function of the Leydig cell system of the testis. The decreased fructose level as well as an increased accumulation of heavy metals in the testis observed in this study is consistent with the earlier report. Scientific evidence suggests that hypofunction of these accessory sex organs affects sperm motility and sperm chromatin stability which may produce infertility (41). The seminal fructose concentration is an assessment of the seminal vesicle and could also give useful indications of the male reproductive function (42).

The level of cellular adenosine triphosphate (ATP) during anaerobic conditions had been widely assessed using LDH activity because it is a stable enzyme. LDH is an oxidoreductase enzyme that catalysis the interconversion of pyruvate and lactate. Cells release LDH into the bloodstream after tissue injury. Animals exposed to the dumpsite had a significant depletion of LDH activity in the serum. Inhibition of LDH which is a key enzyme of glycolytic pathway caused by the dumpsite exposure in the absence of oxygen which is needed for ATP production would slow down the metabolic pathway in charge of energy production. This outcome supports (43,44) who had previously reported that patients with abnormal spermatogenesis had low levels of ATP. Additionally, ATP is necessary for the synthesis of cholesterol, ketone bodies and fatty acids. In aerobic conditions, pyruvate is converted to acetyl CoA catalyzed by pyruvate dehydrogenase which may be oxidized in the tricarboxylic acid (TCA) cycle to produce ATP. The findings in this study thus suggest that heavy metals in the dumpsite could disrupt ATP production by inhibiting pyruvate dehydrogenase when competing with the phosphate group thus uncoupling oxidative phosphorylation. This inhibits energy-linked decrease of NAD+, mitochondrial respiration, and ATP synthesis in the serum. It is interesting to note that a different interplay of events occurred in the testis of the DSE animals, as there was an increase in the level of LDH activity in the testes. Lactate dehydrogenase is the functional marker of ATP and it is required in its production. LDH is also connected with existence and development of germ cells. The increase in testicular LDH activity thus observed in this study may indicate an adaptive mechanism by the testes to assuage germ cell death and azoospermia in DSE rats.

The significant correlation between the testicular LDH, mercury and copper, as well as fructose and nickel and lead in the DSE animals further buttress the fact that the heavy metal bioaccumulation in the testis affected the energy supply to the testis thus the reason for the adaptive mechanism. Polychlorinated biphenyls (PCBs) are one of the groups of persistent organic pollutants and they have been linked to reproductive difficulties (45). In this study PCBs were absent in the testes of the DSE rats. According to a study (46), the PCB levels in the AWSD topsoil were below detection limit which would have been the reason it was not detected in the testis of the DSE rats. This however does not rule out the possibility of the presence of organic pollutants at the dumpsite as there was an accumulation of polyaromatic hydrocarbons (PAHs) according to their study.

Conclusions

The direct exposure to a municipal dumpsite elicits detrimental effects on the energy supplies needed for male reproductive activity. The exposure compromised testicular LDH and decreased the LDH activity in the serum which are markers of cellular ATP. The inhibition of LDH activity and a decrease in fructose levels observed could be possible mechanisms by which direct dumpsite exposure elicited energy depletion and toxicity in the male rat. PCBs were absent thus the biotoxic effects and energy depletion could be linked to heavy metal interactions which was additive, synergistic, antagonistic and individual in nature. Indeed, an alteration in LDH and fructose levels would be expected to compromise the energy supplies to germ cells. The human male has relatively low fertility potential compared with other mammals and is much more susceptible to metal toxicity, there is therefore the need for a critical assessment and discouragement of siting residential areas near dumpsites.

Conflicts of interest

The authors declare they do not have any conflicts of interest.

References

1. Porta D, Milani S, Lazzarino AI, Perucci CA, Forastiere F. Systemic review of epidemiological studies on effects associated with management of solid waste. Environ. Health. 2009; 8(60):1-14.
2. Odewabi AO, Ogundahunsi OA, Odewabi AA, Oritogun KS, Ekor M. Adenosine deaminase activity and immunoglobulin levels as potential systemic biomonitors of occupational hazards and health status in municipal solid waste management workers. Environ. Toxicol. Pharmacol. 2013; 35:1-12.
3. Abul S. Environmental and health impact of solid waste disposal at Mangwaneni dumpsite in Manzini: Swaziland. J. Sustainb. Dev. Afr. 2010; 12 (7):64-78.

4. Aitken RJ, Ryan AL, Curry BJ, Baker MA. Multiple forms of redox activity in populations of human spermatozoa. Mol. Hum. Reprod. 2004; 9: 645-661.
5. Oluyemi EA, Feuyit G, Oyekunle JAO, Ogunfowokan OA. Seasonal variations in heavy metal concentrations in soil and some selected crops at a landfill in Nigeria. Afr. J. Environ. Sci. Technol. 2008; 2: 89-96.
6. Wang Y, Fang J, Leonrd SS, Rao KM. Cadmium inhibits the electron transfer chain and induces reactive oxygen species. Free Radic. Biol. Med. 2004; 36:1434-1443.
7. Chakrabarti SK, Bai C, Subramanian KS. DNA-protein cross links induced by nickel compounds in isolated rat lymphocytes role of reactive oxygen species and specific amino acids. Toxicol. Appl. Pharmacol. 2001; 170(3):153-165.
8. Chowdhury AR: Recent advances in heavy metals induced effect on male reproductive tract toxicity. In: Endocrine and hormonal toxicology. PW Harvey, KC Rush and A Cockburn (eds.) Wiley, Chichester. pp. 355-405,2009.
9. Sikka SC, Wang R. Endocrine disruptors and estrogenic effects on male reproductive axis. Asian J. Androl. 2008; 10(1):134–145.
10. Pizent A, Tariba B, Živković T. Reproductive toxicity of metals in men. Arch. Hig. Rada. Toksikol. 2012; 63(1): 35-46.
11. Li H, Chen Q, Li S,*et al.* Effect of Cr (VI) exposure on sperm quality; human and animal studies. Ann. Occup. Hyg. 2001; 45:505-511.
12. Kumar S, Sathwara NG, Gautam AK, *et al.* Semen quality of industrial workers occupationally exposed to chromium. J. Occup. Health. 2005; 47: 424-430.
13. Yan F, Chen Y, Zuo Z, Chen Y, Yang Z, Wang CH. Effects of tributyltin on epididymal function and sperm maturation in mice. Environ. Toxicol. Pharmacol. 2009; 28(1): 19–24.
14. O'Flaherty CM, Beorlegui NB, Beconi MT. Lactate dehydrogenase-C4 is involved in heparin- and NADH dependent bovine sperm capacitation. Andrologia, 2002; 34 (2): 91-97.
15. Tsujii T, Kamai T, Moriguchi H, *et al.* Seminal lactate dehyrogenase C4 (LDH-C4) isozyme activity in infertile men. Hinyokika Kiyo. 2002; 48 (4): 193–197.
16. Duan C, Goldgerg E. Inhibition of lactate dehydrogenase C4 (LDH-C4) blocks capacitation of mouse sperm *in vitro.* Cytogenet. Genome. Res. 2003; 103 (3–4): 352–359.
17. Dias TR, Martins AD, Reis VP, Socorro S, Silva BM, Alves MG, Oliveira PF. Glucose transport and metabolism in Sertoli cell: relevance for male fertility. Curr. Chem. Biol. 2013; 7: 282-293.
18. Sharifah Norkhadijah SI, Nurizzati Y, Sarva Mangala P, Abidin EZ, Dora EJ. Association between living near to a non-sanitary landfill fingernail cadmium level and health symptoms among children in Nilai. Int. J. Pub. Health. Clin. Sci. 2015;2(3):105-121.
19. Bortey-Sam N, Nakayama SMM, Ikenaka Y, Akoto O, Baidoo E, Mizukawa H, Ishizuka M. Heavy metals and metalloid accumulation in livers and kidneys of wild rats around gold-mining communities in Tarkwa Ghana. J.Environ. Chem..Ecotoxicol. 2016; 8(7):58-68.
20. Adedara IA, Oyebiyi OO, Lawal TA, Adesina AA, Farombi EO. Involvement of oxidative stress in municipal landfill leachate-induced toxicity in boar sperm. Environ. Toxicol. Pharmacol. 2013; 36: 972-978.
21. Adedara IA, Lawal TA, Adesina AA, Oyebiyi OO, Ebokaiwe AP, Farombi EO. Sperm functional parameters and erythrocytes oxidant-antioxidant imbalance during municipal landfill leachate treatment withdrawal in rats. Environ. Toxicol. Pharmacol. 2014; 37:460-467.
22. Adedara IA, Awogbindin IO, Adesina AA, Oyebiyi OO, Lawal TA, Farombi EO. Municipal landfill leachate-induced testicular oxidative damage is associated with biometal accumulation and endocrine disruption in rats. Arch. Environ. Contam. Toxicol. 2015; 68:74-82.
23. Akintunde JK, Oboh G, Akindahunsi AA. Testicular membrane lipid damage by complex mixture of leachate from municipal battery recycling site as indication of idiopathic male infertility in rat. J. Interdiscip. Toxicol.2013a; 6(4): 192-197.
24. Akintunde JK, Oboh G, Akindahunsi AA. Inhibition of key markers linked with spermatogenesis and cellular ATP by subchronic exposure to leachate in a rat model. Arch. Environ. Contam. Toxicol. 2014; https://doiorg/101007/s00244-014-0068-9.
25. Waste Atlas. The World's Biggest Dumpsites. D waste 2014.
26. Luvizutto JFL, Solano M De LM, Martinez MF, Fernandez CD, Umbuzeiro G de A, Camargo JLde. Potential androgenic effects of urban sewage sludge in male rats. Endocrine Disruptors. 2015; 3: 1 e1066656.
27. Farombi EO Adewolo OA& Ajimoko YR Biomarkers of oxidative stress and heavy metal levels as indicators of environmental pollution in African catfish (Clarias gariepinus) from Nigerian Ogun river Int J Environ Res Public Health 4158-165 (2007)
28. Adeyemo A, Morenikeji OA. Helminths and heavy metals in soils from a dumpsite in Ibadan city. Nigeria. J. Prev. Med. Hyg. 2017; 58: E328-E333.

29. Telišman S. Interactions of essential and/or toxic metals and metalloid regarding interindividual differences in susceptibility to various toxicants and chronic diseases in men. Arh. Hig. Rada. Toksikol. 1995; 46:459-476.
30. Sharpe RM, McKinnell C, Kivlin C, Fisher JS. Proliferation and functional maturation of Sertoli cells and their relevance to disorders of testis function in adulthood. Reproduction 2003; 125:769-784.
31. Rosner M, Dolznig H, Schipany K, Mikula M, Brandau O. Human amniotic fluid stem cells as a model for functional studies of genes involved in human genetic diseases or oncogenesis. Oncotarget. 2010; 2:705–712.
32. ATSDR. Agency for Toxic Substances and Disease Registry Minimum risk levels (MRLs). 2013. Available online at http://wwwatsdrcdcgov/mrls/pdfs/atsdr_mrls_february_2012pdf
33. Klaassen C, Watkins J. In: Casarett and Doull's essentials of toxicology. McGraw-Hill. pp.512,2003.
34. World Health Organisation International year of fresh water. General Assembly Resolution. 2008; A/RES/55/196 wwwwater year 2003org Accessed 10 Feb 2019.
35. de Souza Predes FM, Diamante AS, Dolder H. Testis response to low doses of cadmium in wistar rats. Int. J. Exp. Pathol. 2010; 91 (2):125–131.
36. Lanning LL, Creasy DM, Chapin RE. Recommended approaches for the evaluation of testicular and epididymal toxicity. Toxicol. Pathol. 2002; 30(4): 507–520.
37. Ohtani K, Yanagiba Y, Ashimori A, *et al.* Influence of injection timing on severity of cadmium-induced testicular toxicity in mice. J. Toxicol. Sci. 2013; 38(1):145–150.
38. Hess RA. Effects of environmental toxicants on the efferent ducts epididymis and fertility. J. Reprod. Fertil. 1998; 53: 247-259.
39. Apostoli P, Catalani S. Metal ions affecting reproduction and development. Met. Ions. Life. Sci. 201; 8:263-303.
40. Martinez CS, Torres JG, Peçanha FM *et al.* 60-Day Chronic Exposure to Low Concentrations of $HgCl_2$Impairs Sperm Quality: Hormonal Imbalance and Oxidative Stress as Potential Routes for Reproductive Dysfunction in Rats. PLoS One 2014; 4: 9(11).
41. Gonzales GF, Villena A. Influence of low corrected seminal fructose levels on sperm chromatin stability in the semen from men attending an infertility service. Fertil. Steril. 1997; 67:763.
42. Trang NT, Sang TT, Hoang N, Khanh NTG, Duc TT. Assessment of the level of seminal zinc and fructose concentration in seminal plasma of Vietnamese infertile men. MOJ Biorg. Org. Chem. 2018; 2(4):185-190.
43. Parlakpinar H, Tasdemir S, Polat A, *et al.* Protective effect of chelerythrine on gentamicin-induced nephrotoxicity. Cell Biochem. Funct. 2006; 24:41–48.
44. Walker PDY, Barri SV. Oxidant mechanisms in gentamicin nephrotoxicity. J. Ren. Fail. 1999; 21:433.
45. USEPA. Ambient water quality criteria for chloride. Washington DC: US Environmental Protection Agency Criteria and Standards Division 1998.
46. Oketola AA, Akpotu SO. Assessment of solid waste and dumpsite leachate and topsoil. Chem. Ecol. 2014; 31(2):134-146.